

LA-UR-12-26089

Approved for public release; distribution is unlimited.

Title: SPE Near Field Velocity Data Corrections and Analysis

Author(s): Steedman, David W.

Intended for: Distribution of analysis results to project participants




Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

SPE Near Field Velocity Data Corrections and Analysis


David Steedman/LANL
22 October 2012

Outline



- **Gage Corrections**
 - General
 - Methodology
- **Review of data**
 - SPE-1
 - SPE-2 and SPE-3
- **Summary**
- **Recommendations**

Gage Corrections




- **Gages in holes 1 through 6 are assumed to have rotated**
 - The exception is level 3 (*i.e.*, 50-ft depth) gages
 - Jeff Thomsen/ARA determined rotation angle from relative R and T component accelerations
- **All gages in holes 7 through 11 were assumed to have not rotated**
- **All corrected data have been posted to the UNR site**
 - `mirror1/home/steedmand/speNFcorrections`
- **For the bulk of this presentation we note that prior analysis has focused on record peak values and arrival times**
 - With corrected data we can examine waveforms
 - We review waveforms for level 1 (180-ft) and level 2 (150-ft) depths

Comments on Methodology


- One hypothesis prescribed that joint sets at SPE site “turned” the velocity vector away from a true radial direction
- With recovery of SPE-3 data from newly installed accelerometers the discussion properly focused on shear wave effects (September ‘12)
- Why revisit this now?
 - Xu email during data correction activity (10/5/12):
 - 5) The new gages in SPE3 have significant tangential components, **causing ~12 degrees deviation of the velocity vector away from the spherical radial direction** (a mean of 0.23 for tangential/radial ratio in SPE 3 corresponds to ~12 degrees). The rotation angles calculated should contain this type of error information.
 - This ignores the consensus that the transverse magnitudes are high due to shear wave effects (\neq deviation of vector)

“Turning Vector” vs. Shear Wave Arrivals





- Turning of the vector implies a realignment of the shock front relative to test geometry

- Off-radial components would be of *similar* shape to radial but with reduced amplitude
- Specifically, peaks would be coincident




- But in most cases V_P^T lags V_P^R by a significant delay
- Also $V_P^T(t)$ is characteristically different from V_P^R
- History is characteristic of the slower shear wave
 - See data from records that did not require correction such as SPE-3 canister 7-2

Other Examples




SPE-3 8-2



V^T at $t_P^R \cong 0$ (not 23%)



SPE-2 4-1
(Corrected)

Study Full Data Set




- Look at SPE-1, then the SPE-2/SPE-3 pair
 - Different scaled ranges to gages
 - Different yields
- In following plots:
 - Radial is blue
 - Transverse is red
 - Longitudinal is green
 - All time scales are identical
 - Amplitude is self-scaled

SPE-1 Data– 10-m Range



150-ft Depth




Insufficient Data to Estimate Rotation in 3-1

180-ft Depth



Hole 2

Hole 3


SPE-1 Data– 10-m Range, 150-ft depth (45° from source)



- R-component and L-component should be equal
 - Shear arrivals alter L-component at both locations
- Shear arrivals at two different azimuths are coincident in time
 - Suggests a common source
 - *i.e.*, the explosive source vs. random joints






Hole 2





Hole 3

SPE-1 Data– 20-m Range




150-ft Depth




180-ft Depth

SPE-1 Arrival Times

- Shear speed about equal to loading speed
 - Times to peak and peak amplitudes are likely obscured by shear wave
- Tight trend in shear arrivals suggest a common source




SPE-2 and SPE-3 Data Repeatability





- **Several methods indicate that SPE-2 created a damaged medium for SPE-3**
 - Near field late arrivals and lower amplitude
 - Cross-hole survey
 - Infrasound
 - Slant hole fractured core
- **But gage-by-gage SPE-3 data are remarkably similar to SPE-2 data**
 - Most notable for R-components
 - But also some T- and L- components


SPE-2/SPE-3– 10-m Range, at Shot Depth



- No 3-2-R in SPE-3, but nearby 8-2-R in SPE-3 is very similar to 3-2-R in SPE-2






SPE-2




SPE-3

SPE-2/SPE-3– 20-m Range, Shot Depth






SPE-2





SPE-3

SPE-2/SPE-3– Hole 2



SPE-2







SPE-3




150-ft Depth

180-ft Depth

SPE-2/SPE-3– Three-component Repeatability



SPE-2



SPE-3



Hole 2

Hole 4

Hole 5


SPE-3– Revisit Can 9-2

- 9-2-R anomalously high
- Review all SPE-3 at 20-m range and shot depth
- Strong 9-2-T shear coincides with 9-2-R inflection
 - Amplitude increases by 60% after inflection
 - That is of the order increase that was experienced by SPE-1 can 2-2-L after shear arrival




SPE-2 Arrival Times

- Similar characteristics as for SPE-1; i.e.,
 - Shear arrivals “interfere” with peaks
 - Trend in shear arrivals suggests a common source



SPE-3 Arrival Times


- Similar characteristics as for SPE-1 and SPE-2



SPE-2/SPE-3– Summary


- **SPE-3 histories resemble SPE-2, even down to small perturbations**
- **Similarities/Differences**
 - Explosive sources for these events were nearly identical
 - The rock medium for SPE-3 was damaged by SPE-2
- **It follows that the consistent waveforms reflect consistent source characteristics**

Summary




- **There appear to be significant shear waves in the SPE ground shock environment**
- **Arrival times suggest a point source (explosive) rather than a distributed source (rock joints)**
- **Character of waveforms is similar when considering like range and depth across differing azimuths**
- **Character of waveforms is repeatable across experiments with repeated source yield and location**

Discussion




- **Would a spherical source cause similar shear waves?**
- **If so, do we want to study cylindrical sources or spherical sources?**


Theoretical Radiation from a Cylindrical Source




- From a moment tensor for cylindrical charge
 - $L/D = 4$
 - Significant SV radiation predicted



P-Radiation



SV-Radiation




SH-Radiation

Recommendation

- If we are interested in cylindrical sources
 - Proceed as planned
- If we are interested in spherical sources
 - Numerical simulations are required to compare environments created by cylindrical sources to those from spherical sources
 - Design SPE-4 as a spherical source
 - If study/SPE-4 show that there is no difference → Proceed as planned
 - If study/SPE-4 show that there is a difference → Reconsider future event designs

Addendum




- **Discussion: how are different sources of non-radial motion (turning shock vector vs. shear arrivals) manifest in the recorded data?**

Comments on Phenomenology


- It has been hypothesized that joint sets at SPE site turn the velocity vector away from a true radial direction
 - This was based on observing relatively large transverse motions
 - But for SPE-1 and SPE-2, often the transverse amplitude was *greater* than the radial; by more than factor of 5 in some cases
 - No credible explanation provides for this large a variation
 - Moreover, the site joints are 3-D and there is no evidence suggesting the joints have similarly affected the longitudinal (*i.e.*, vertical) measurements
- SPE-3 data from newly installed accelerometers do not display these excessive transverse amplitudes
- Why revisit this now?
 - Xu email during data correction activity (10/5/12):
 - 5) The new gages in SPE3 have significant tangential components, **causing ~12 degrees deviation of the velocity vector away from the spherical radial direction** (a mean of 0.23 for tangential/radial ratio in SPE 3 corresponds to ~12 degrees). The rotation angles calculated should contain this type of error information.

“Turning Vector” vs. Shear Wave Arrivals




- Turning of the vector implies a realignment of the shock front relative to test geometry
- If this were true:
 - Off-radial components would be of *similar* shape to radial but with reduced amplitude
 - Peaks would be coincident, or nearly coincident, and waveforms would be similar (duration, shape, *etc.*)
- This result would be indistinguishable from a rotated gage package except for one point:
 - The site joint systems are 3-D, consequently, the phenomenon affecting the transverse motion should occur to a similar degree on the longitudinal measurements
 - Conversely, rotation of the gage package would have no effect on a longitudinal history


“Turning Vector” vs. Shear Wave Arrivals (cont.)




- The uncorrected histories from SPE-1 can 1-1 provide an example
 - The **radial** and **transverse** waveforms are nearly identical except for lower transverse peak and a post-shock negative drift in the radial
 - But the **longitudinal** history is radically different, and has a peak about 20% of the radial peak
 - This combination of effects suggests that the high transverse is due to a rotated gage, not a rotated shock front




“Turning Vector” vs. Shear Wave Arrivals



- We can contrast this with a can where there is high confidence that it did not rotate
- The peak **transverse** velocity (V_p^T) lags the peak **radial** velocity (V_p^R) by a significant delay
- The **transverse** history ($V_p^T(t)$) is characteristically different from the radial history (V_p^R)
- The **longitudinal** history ($V_p^L(t)$) is also characteristically different from the radial history (V_p^R) with a small peak






Review all Gages in Holes 7 – 11 (Except for Slant Hole 10)



- To study this further, review SPE-3 data for all data that did not require rotation
 - Transverse peak value relative to radial peak
 - Time of transverse peak relative to radial peak
 - Value of transverse velocity at time of radial peak
 - Review longitudinal response as well
 - In particular, review longitudinal histories at the shot depth (level 2)
- Focus on level 1 and level 2 gages
 - Ground surface, weathered-to-intact interface and water table complicate the level 3 response
 - We will examine levels 1 through 4 for hole 9

Hole 7






There is no characteristic similarity between radial and transverse; although after 12 ms the respective curves are nearly the same

7-1

7-2

Hole 8








There is no characteristic similarity
between radial and transverse

Transverse has $\frac{1}{2}$ the first
phase duration as radial




8-1

8-2

Hole 9



Hole 11




There is no similarity
between radial and transverse

Resolved Horizontals

- Take the peak radial velocity
- Use the Transverse velocity value at the time of the peak radial
- Compute the peak of the resolved vector in the horizontal plane (i.e., neglect longitudinal contribution)
- The transverse component provides an insignificant contribution to the resolved horizontal velocity
- The vector is not deviating from the geometric radial


Can	Peak Radial (m/s)	Transverse at Time of Peak Radial (m/s)	Resolved peak (m/s)
7-1	1.054	0.0271	1.054
7-2	1.255	0.157	1.265
8-1	0.98	0.189	0.998
8-2	5.249	0.537	5.276
9-1	1.119	0.225	1.141
9-2	5.205	0.691	5.251
9-3	1.565	0.259	1.586
9-4	0.898	0.005	0.898
11-1	0.604	0.017	0.604

Hole 7-11 Observations



- **Hole 7**
 - **Level 1**
 - The transverse peak lags the radial peak by a considerable time
 - The transverse amplitude at the time of radial peak is zero
 - The general character of the radial and transverse histories are quite dissimilar, although they nearly overlay after 12 ms time
 - **Level 2**
 - The transverse peaks lags the radial peak by a considerable time
 - The transverse amplitude at the time of radial peak is about 12% of the radial peak
 - The transverse amplitude at the time of the radial peak is about 1/3 of the transverse peak
 - The general character of the radial and transverse histories are quite dissimilar, although they nearly overlay after 12 ms time

Hole 7-11 Observations (cont.)



- **Hole 8**


- **Level 1**

- The transverse peak, while about $\frac{1}{2}$ the radial peak amplitude, lags the radial peak by about 1 ms
 - The transverse amplitude at the time of radial peak is about 10% of the radial peak, and is less than $\frac{1}{2}$ the value of its own peak
 - The general character of the radial and transverse histories are quite dissimilar

- **Level 2**

- The transverse peak, while about 20% of the radial peak amplitude, lags the radial peak by about 1 ms
 - The transverse amplitude at the time of radial peak is about 10% of the radial peak, and is less than $\frac{1}{2}$ the value of its own peak
 - The general character of the radial and transverse histories are quite dissimilar; with the first phase of the transverse being about one half the duration of the radial outward phase

Hole 7-11 Observations (cont.)



- **Hole 9**


- **Level 1**

- The transverse peak, while about 30% of the radial peak amplitude, lags the radial peak by about 1 ms
 - The transverse amplitude at the time of radial peak is about 15% of the radial peak, and is about 30% of the value of its own peak
 - The general character of the radial and transverse histories are not similar

- **Level 2**

- The transverse peak, while about 17% of the radial peak amplitude, lags the radial peak by about 1 ms
 - The transverse amplitude at the time of radial peak is about 12% of the radial peak
 - The general character of the radial and transverse histories are similar

Hole 7-11 Observations (cont.)



- **Hole 9 (cont.)**


- **Level 3**

- Except for sign, the transverse history is characteristically similar to the radial history
 - The radial and the transverse peaks are nearly simultaneous
 - The transverse peak amplitude is about 20% of the radial peak amplitude

- **Level 4**


- The transverse peak, while about 20% of the radial peak amplitude, lags the radial peak by about 2 ms
 - The transverse amplitude at the time of radial peak is 0.
 - The general character of the radial and transverse histories are quite dissimilar

Hole 7-11 Observations (cont.)



- **Hole 11**
 - **Level 1**
 - The transverse peak lags the radial peak by about 4 ms
 - The transverse amplitude at the time of radial peak is nearly 0.
 - The transverse peak amplitude is 8% of the radial peak amplitude
 - The general character of the radial and transverse histories are quite dissimilar

Hole 7-11 Observations (concl.)



- **For all canisters**
 - **Peaks**
 - The transverse peak generally lags the radial peak by several ms
 - The amplitude of the transverse waveform at the time of radial peak is generally less than 10% of the radial peak (close to 0. in some cases)
 - The longitudinal peaks are generally less than 10% of the radial peaks
 - **Character**
 - The character of the transverse waveform is generally considerably different from the character of the radial waveform
 - In nearly all cases the character of the longitudinal waveform is considerably different than the character of the radial waveform
- **These characteristics all argue for late time phenomena that are not concurrent with the main shock**
 - The “high” (i.e., non-zero) transverse measurements are not indicative of a turning shock vector